References

Abbas. G. Khattak.A, Mir. A, Ishaque.M, Hussain.M, Wahedi.M, Ahmed. MS, 2012, Effect of organic manures with recommended dose of npk on the performance of wheat, Journal of Animal plant Science, P.no. 683-387.

Abdo.C, Nemer.N,Nemer.G, Abou.Y, Atamian.H, Kawer.N, 2006, Isolation of Beauveria species from Lebanon and evalution of its efficacy against the ceder web-spining sawfly, Cephalcia tannourinensis, BioControl, V.no53, P.no. 341-352.

Ahmad .A, Ahmad.I, Khan.MS, 2006, Screening of free living rhizospheric bactria for their multiple plant growth promoting activities, Department of Agriculture, Aligarh.

Alves SB, Perera RM, 1989, Production of metrazium anisopliae and beauvria bassiana in plastic trays, 14, p.n. 188-192.

Babalola.OO, 2010, Beneficial bacteria of agriculture importance, Biotechnology letters, V.no.32, P.no.1559-1570.

Bhat .JM, Alagawadi. AR, 1998, Seed borne nature of Azotobactor Chrocoocum in chili and its role in seed germination and plant growth, Hindustan Antibiotic Bull, P.no. 20-30.

Benitez.T, Rincon.AM, Limon.MC, Codon .AC., 2004, Biocontrol mechanism of Trichoderma strains, Environmental Microbiology , V.no.1, P.no. 249-60

Brewin NJ,2002, Pods and nods: A now look at symbiotic nitrogen fixing bacteria, P.no 113-11.

Baset.MA, Shamsuddin. ZH, 2013, Rhizobium as a crop enhancer and Biofertilizer for increased cereal production, African Journal of Biotechnology, P.no.9.

Batta.YA, 2007, Biocontrol of Almond bark beetle (Scolytus amygdale geunin- meneville, Coleoptera scolytidae) using Beauveria.bassiana. Journal of Application Microbiology, V.no.103, P.no.1406-1414.

Chandler.D, Davidson.G, Prant.W, Graves.J, Tatchell.GM, 2008, Microbial biopesticides for integrated crop management an assessment of environment and regulatory sustainability, University of Warwick, V.no19. P.no.275-283.

Chet.I, Inhar, Hehrew, jul1994, Biological control of fungal pathogens, Application of Biochemistry biotechnology, V.no.1, P.no. 37-43.

Cloyd .R,1999, The Entomopathogenic fungus *Verticillium lecanii*, University of Illinois.

Cloyd .R,1999, The Entomopathogenic fungus *Metragium anisoplia*, University of Illinois.

Denis.HB, 1998, Formulation of mycoinsecticides, Formulation and microbial pesticides, P.no.131-185.

Donald.WR, Hajek.AE, 1992, Entomopathogenic fungi as bioinsecticides, Frontiers in industrial mucology, P.no.144-159.

Edson.H, Pedr.M, Joao.A, Luis.H, Martin.CH, Peralta and Alcides.M, 2001,Effect of Biofertilizer and neem oil on the entomopathogenic fungi Beauveria bassiana and Metrazium anisoplia, Braz arch boil technol, V.no. 4.

Gamal.ME, Ghazal and, Shahhat.M, 2012, Psiological and Phytochemical response of *Foeniculum vulgare var. vulgare* Mill and *Foeniculum Vulgore var, azoricum* Mill to Bio-organic Manuras partial or full substitute for Inorganic Amendment, Austraian Journal of Basic and Applied Science, P.no.266-277.

Gilardi.G, Baudino.M, Gullino.MC, Garibaldi.A, 2008, Attempts to control fusarium root rot of bean by seed dressing. Common Agriculture Application Biology, P.no. 75-80.

Goettel MS, Koike M, Kim.J, Aiuch.D Shinya R and Brodeu. J , 2008, Potential Lecanicillium spp for management of insect nematodes and plant diseases, Journal of invertebrates pathogen, 98, 256-261.

Hafte.M,, Altier. N, 2010, Fluorescent Pseudomonas as biocontrol agents for sustainable agriculture system, Res. Microbiology, P.no.464-471.

Jin. SF, Feng MG, Chen JQ, 2008, Institute of Microbiology, China, Pest Management Science, V.no.10, P.no.1008-1014.

Jos.M, Raaijmakers.T, Poulitz.C, Stainberg.C, Alabouvette.C, Loccoz.YM, 2009, The Rhizosphere: a playground and batter field for soil born pathogen and beneficial microorganism, V.no.321, P.no.341-361.

Kamp AM, Bidochka.MJ, 2002, Conidium production by insect pathogenic fungi on commercial available agar,

Kaur K, Goyal.S, Kapoor.KK, 2008, Impact of organic fertilizer with and without chemical fertilizer on soil chemical properties and the establishment of nitrogen fixing bacteria in the Rhizosphere, Page no.313-316.

Kethr.FB, Samia.A, Dhouh.S, Majda.D, Jihane.C, Kcouthan.L, Mohamed.M, Ahmed.D, Zine.M, 2008, Chemical composition antibacterial and antifungus activity of Trichoderma species growing in Tunisia, Annals of Microbiology, V.no.58, P.no.303-308.

Khetar SK, 2001, Microbial pesticides, Marcel Dekker Ins, Newyork p.n. 300.

Khahid.A, Arshad.M, and Zahir.ZA, 2004, Screening plant growth promoting rhizobacteria for improving growth and yield of wheat, Journal of Applied Microbiology, V.no.96, P.no. 473-480.

Khan.MS, Ahmad.F, Zaidi.A, Oves.M, 2013, Functional aspect of Phasphate solubilizing bacteria: Importance in crop production, Bacteria in Agronimy: crop productivity, P.no.237-263.

Knudsen.GR, Eschen.DJ, Dandurand. LM and Wang. Z, Method to enhancer growth and sporulation of pelletized biocontrol fungi, University of Idaho.

Kumar .V, Singj. KP, 2001, Enriching vermicompost by nitrogen fixing and phosphate solubiliging bacteria, Bioresour Technol, P.no.173-175.

Leinhos. V, Vacek.O, 1994, Biosynthesis of auxin by phosphate solubilizing rhizobacteria from wheat, Institute of microbiology, Germany, V.no.149, P.no31-35.

Leinhos.V, 1994, Effect of Ph and glucose on auxin production of phosphate solubilizing rhizobacteria in vitro, Microbiology research, V.no.149, P.no.135-138.

Lomer CJ, Priore C, Kooyman G, Goettel MS, Johnson DL,1997, Development of Metrazium species for the control of grass hoper and locusts. Microbial control of grasshopper and locusts, 265-286.

Marino.S, Cecilia.M, Critina.C, 2010, A comparison of preservation method for Trichoderma harzianum cultures, Rev Iberoam Miccol.

Marilyn.J, Roossinck, 2008, Benificial microbes for Agriculture, The Samuel Roberts Nobel Foundation.

Mia.MA, Shamsuddin.ZH, Watab.Z, Mahmood.M, 2007, Associative nitrogen fixation by Azospirillum and Bacillus species in Banana, V.no.16, P.no.11-15.

Montesinos .E, sep.2003, Develoment, registration and commercialization of microbial pesticides for plant protection, Environmental Microbiology, P.no.245-252.

Moore D, 2008, A plague on locust, The Lubilosa Story, Outlook pest management 19, 14-117.

Nezarat.S, Gholami.A, 2009, screening plant growth promoting rhizobacteria Pak. J. Biol. Sci., P.no. 26-32.

Nadeau MP, Dunphy GB, and Boisvert JL, 1996, Development of Erynia conidial on the cuticle of the adult black flies simulium rostrutum and simulium decorum. Journal of invertebrate pathogen, 50-58.

Olabiyi.T.L, Ruoccom.M and Leenzuize.S, 2014, Pathogenic fungi and bio-control agents competitive bio-assay research, Building organic bridge, Germany,p.no.733-736.

Patel .B, Phanse. N, A Prrctical Approach Microbiology, P.M.B Gujarati Science College, Indore, p. no. 121-123,127.

Ramazon. C, Figen.D, Adil.A, Fikrettin.S, 2006, Groth promotion of plant by plant growth-promoting Rhizobacteria under green house and two different field soil conditions, Soil biology and Biochemistry, Vol.no.38, P.no.1482-1487.

Relic.E, Penetx, Estrade.G.M, Galinowaski.J, Krishnaa.HB, Puppka.SG, 1994, Nod factor of Rhizhobium area a key to the legume door. V.no.1 P.no.177-178.

Robl.D, Sung L.B, Novakovich.jh, Marangoni Pr, Zawadneak.MA, Dalzoto .PR, Gabardo.j, Pimentel.I, 2009, Spor production in Paecilomyces lilacinus Samson strains on agro-industrial residues. Brag J Microbiol, P.no 296-300.

Sajda MU, Zaki FA, Jan MA, wanir A, 2007, Comparative susceptibility of different Grup stages to local strain of beavurea bassiana. Journal of entomopathogenic, 4, p. no.56-59.

Sansinenea. E, Ortiz A, 8 nov.2014, Photoprtection for Bacillus thuriengenesis based biopesticides, Biotecnology letter.

Sawada.H, Kuykendall. LD, Young.JM, 2003, changing concepts in the systematic of bacterial nitrogen fixing legume symbiont, General of application biology, V.no.3 P.no.155-179.

Singh. S, Chand. H, 2006, Screening of bioagent against root rot of mung bean caused by Rhizoctonia soloni,Common Agriculture Application Biology, P.no. 33-35.

Sharma.K, Dak.G, Agrawal.A, Bhatnager.M, Sharma.R. 2007, Effect of phosphatsolubilizing bacteria on the germination of cicer arietinum seeds and seedling growth, Department of Botany, Udaipur.

Sharma.p, Verma.V, Ramesh.R, Saravanan SD, Deep.S, Mahesh.S, Dinesh.S, 2014, Biocontrol genes from Trichoderma species: Areview, African Journal of Biotechnology, V.no.10

Soesanto.S, Mugiastuti, Mana, Abdul.M, Wachjadi.M,2013, Ability test of several antagonistic to control potato bacterial wilt in the field, Jendang soedirman University of Indonesia.

Stephen RF, Travis RG, 2014, I nsecticidal agents and uses thereof, Biotelliga hoolding limited.

Stephen. A, Rehner, Andrew.M, Hosung.G, Jennifer L, Devott.L, Humber.RA, 2011, Phylogeny and systematic of the anamorphic entomopathogenic genes Beauveria bassiana, Mycologia, P.no.1055-1070.

Strasser O, kohlic and Romheld V, 1999, Overestimation of apoplastic root of soil groth plant, 210, 179-187.

Susan G.W, Basinger .JF, Bao.X, Rodrigues R.J, Redman. RS.,2013, Method for increasing plant growth using the fungus Trichoderma harzianum, University of Canada and Washington.

Sytnikov DM, Datsenko.VR, 2007, Efficacy of biological preparation of soyabean root nodules bacterial modified a homoglobous lectin, Prikl Biokhim Microbiology, p.no.304-310.

Triq. MB, Henis.J, Keith.AP, 1999, Microbial Biopesticides, Biopesticides: use and delivery, V.no.5, P.no 23-44.

Travis. RG, Callaghan.MO, 2003, Environmental impacts of Bacterial insecticides progress in biological control, V.no.1, P.no.119-149.

Vidhysekaran. P and Muthamilan.M, 1995, Development of formulation of P.seudomonas fluorescens for control of chick pea wilt, Agriculture University Coimbatore.

Vande sende K, Bisseling.T, 1997, signalling in symbiotic root nodules formation, P.no.127-142.

Viviene .NM, Felix. DD, 2004, Potential use of rhizobium bacteria as promoters of plant growth for increased yield in landraces of African cerea crops, African journal of Biotechnology, V.no.1, P.no.1-7.

Wraigh SP, Inglis GD and Goettel MS, 2007, Field manual of Techniques in invertebrates pathology, 2nd addition, 223-248.

Yang.J, Zang.KQ, 2014, Biological control of plant parasitic nematodes by nematophagous fungus, Fungle diversity Research series, V.no.23, P.no.231-262.

Yoav Bashan, S.R prabhu, J.P Hernandez, 31 oct.. 2013, Advances in plant growth promoting technology formulation and practical perspectives (1998-2013).

Young.T, Min.S, Nakoo.H, Dong.S, 2010, Study on selective media for isolation of Entomopathogenic fungi, International Journal of Industrial Entomology, P.no. 7-12.

Zhou.Y and H, Shen. F, 1998, Applying Beauveria bassiana to control rhytidodera browrigii,journal of Beijing forest university, V.no.4, P.no.65-69.

Zimmermann.G, 2006, the entomopathogenic fungus M. anisoplie and its potential as a biocontrol agent.

.